
Small RNAs – their biogenesis, regulation and function in embryonic stem cells*

Joshua E. Babiarz^{1§} and Robert Blelloch^{1,§}, ¹The Eli and Edythe Broad Center of Regeneration Medicine and Stem Cell Research, Center for Reproductive Sciences, and Department of Urology, University of California, San Francisco, California 94143, USA

Table of Contents

1. Introduction	1
2. The Biogenesis of miRNAs and related small RNAs	2
3. The role of small RNAs in embryonic stem cells	3
4. Transcriptional regulation of the miRNA content of a cell	4
5. Post-transcriptional regulation of the processing of miRNAs	5
6. miRNAs and induced pluripotency	7
7. Non-canonical miRNAs and endogenous siRNAs	8
8. Conclusions	10
9. References	10

1. Introduction

Mammalian development follows a carefully orchestrated unfolding of cell fate decisions whose time, place, and specialization must occur robustly and consistently to end with a healthy adult organism. How this feat is achieved time and time again remains a mystery and is of intense interest to the biological and medical communities. One of the central aspects of this feat is the rapid shifts in molecular programs during developmental transitions. These shifts must occur at multiple levels including the regulation of transcription, mRNA stability, protein translation, protein stability, and protein activity. Regulation occurs with the help of transcription factors (activators and suppressors), epigenetic enzymes, non-coding RNAs, RNA binding proteins, ubiquinating enzymes, kinases, phosphatases, and more. Of great interest are a specific group of non-coding RNAs, the microRNAs, which are only 18–24 nucleotides in length and function by both destabilizing and inhibiting translation of mRNAs. microRNAs are compelling regulators of developmental progression as their promiscuous nature allows a single miRNA to regulate hundreds of mRNAs, simultaneously allowing rapid shifts in the transcriptome and proteome of cells.

*Edited by Kevin Eggan and George Daley. Last revised May 01, 2009. Published May 31, 2009. This chapter should be cited as: Babiarz, J.E., and Blelloch R., Small RNAs – their biogenesis, regulation and function in embryonic stem cells (May 31, 2009), StemBook, ed. The Stem Cell Research Community, StemBook, doi/10.3824/stembook.1.47.1, <http://www.stembook.org>.

Copyright: © 2009 Joshua E. Babiarz and Robert Blelloch. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

§To whom correspondence should be addressed. E-mail: babiarzj@stemcell.ucsf.edu, blelloch@stemcell.ucsf.edu




Figure 1. The Cononical miRNA biogenesis pathway. pri-miRNAs are transcribed primarily as RNA pol II transcripts. The pri-miRNAs are processed co-transcriptionally by the Microprocessor (Drosha/Dgcr8). Following Microprocessor cleavage, Exportin-5 transports the pre-miRNA hairpin into the cytoplasm. There, the pre-miRNA is cleaved by Dicer, resulting in a miRNA/miRNA* complex. With the help of TRBP, the mature miRNA is loaded into the Argonaute subunit of the silencing complex. This complex then goes onto silence target mRNAs post-transcriptionally by translational inhibition and/or transcript destabilization. Adapted from Babiarz et al., 2008.

2. The Biogenesis of miRNAs and related small RNAs

MicroRNAs undergo a series of processing events before maturing into a functional miRNA-silencer complex (Figure 1). Typically, miRNAs are initially expressed as part of a longer RNA polymerase II (pol II) driven transcript, the primary miRNA (pri-miRNA). These longer transcripts can be either coding (often with the miRNA within the introns), or non-coding (Cai et al., 2004; Lee et al., 2004). The miRNAs themselves are contained within hairpin folds along the length of the pri-miRNAs. These hairpins are recognized by the RNA binding protein, Dgcr8 (Denli et al., 2004; Gregory et al., 2004; Han et al., 2004; Han et al., 2006; Landthaler et al., 2004), which then directs the RNase III-containing enzyme, Drosha, to cleave the RNA at the base of the hairpin resulting in the precursor miRNA (pre-miRNA) (Lee et al., 2003). Dgcr8 and Drosha form the Microprocessor complex. Two pieces of evidence suggest that the Microprocessor processes the hairpins as the pri-miRNA is being transcribed, as opposed to following transcription. First, Drosha and RNA pol II can be cross-linked to DNA near the pre-miRNA hairpins, suggesting that the Microprocessor is in close proximity to the DNA (Morlando et al., 2008). Second intronic pre-miRNAs can be released by the Microprocessor prior to splicing of the surrounding exons (Kim and Kim, 2007).

Following production of the pre-miRNA hairpin by the Microprocessor, Exportin-5 transports the pre-miRNA to the cytoplasm (Bohnsack et al., 2004; Lund et al., 2004; Yi et al., 2005; Yi et al., 2003). Once in the cytoplasm, a second RNase III-containing enzyme, Dicer, cleaves the pre-miRNA to liberate the mature miRNA and the complementary “star” sequence (Reviewed in (Hammond, 2005)). Similar to the Microprocessor, where the RNase III Drosha pairs with the RNA-binding protein Dgcr8, Dicer pairs with the human immunodeficiency virus-1 transactivating response RNA-binding protein (TRBP) (Chendrimada et al., 2005; Haase et al., 2005). However, unlike the Microprocessor, where Dgcr8 is essential for activity (Wang et al., 2007), TRBP appears to be dispensable for Dicer activity in vitro, but important for choosing the correct strand for transfer to the silencing complex and downstream gene silencing (Chendrimada et al., 2005; Gregory et al., 2005; Haase et al., 2005).

In addition to miRNA hairpins, Dicer is also capable of cleaving long, double stranded RNAs to small RNAs. The resulting small RNAs are named small interfering RNAs (siRNA). Endogenous siRNAs are found across many species and their function may be even more evolutionary conserved than miRNAs. Indeed, Dicer exists in the absence of the Microprocessor in many species outside of the metazoan lineage (Grimson et al., 2008). siRNAs are capable of directing heterochromatin in non-mammalian eukaryotes, such as *S. pombe* (Verdel and Moazed, 2005), *Tetrahymena*

(Liu et al., 2007) and *Drosophila* (Riddle and Elgin, 2008). Whether there is a similar role for siRNAs in mammalian cells remains unclear. However, endogenous siRNAs have been recently discovered in mammals (Babiarz et al., 2008; Tam et al., 2008; Watanabe et al., 2006; Watanabe et al., 2008; Yang and Kazazian, 2006).

Following processing by Dicer, one strand of the duplex miRNA enters into the silencing complex, the RNA-induced silencing complex (RISC). RISC contains the Argonaute proteins and plays a central role in executing miRNA based silencing. The mechanism for choice of strand and removal of the opposing strand is not entirely clear. However, it appears the energetically less stable 5' end of the duplex is preferentially loaded (Khvorova et al., 2003; Schwarz et al., 2003), while the passenger strand is cleaved by Argonaute-2 (Ago2) (Rand et al., 2005). Ago2 is one of four argonaute proteins (Ago1 to 4) in mammals (Carmell et al., 2002). However, it is only one with cleavage (or “slicer”) activity (Liu et al., 2004; Meister et al., 2004; Rand et al., 2004) and is the only one essential for development (Alish et al., 2007; Liu et al., 2004; Morita et al., 2007). Although Ago2 is essential for normal development, it appears that the four Argonautes may functionally overlap in miRNA silencing. Deletion of all four Argonautes in mouse ES cells results in a loss of silencing and apoptosis (Su et al., 2009). Reintroduction of any of the four Argonautes in these cells results in a rescue of both post-transcriptional silencing and apoptosis (Su et al., 2009).

The miRNA loaded RISC decreases protein levels encoded by the target mRNAs, both by destabilizing the mRNA and by inhibiting translation. Although the precise mechanism for how a miRNA finds a particular target and inhibits translation is the subject of intense investigation, many rules are already known (Bartel, 2009). The major determinant for miRNA targeting is the “seed” sequence – the 7mer sequence present at positions 2 through 8 of the mature miRNA (Lewis et al., 2003). However, there are other components of the miRNA sequence that appear to influence target repression, including nucleotides 12 to 17 in the miRNA and the position of the target site in the 3' UTR of the target gene (Grimson et al., 2007). Roughly one-third of human genes contain conserved miRNA binding-sites in their 3' UTR (Lewis et al., 2005), and the number of recognized conserved sites has recently increased with more sensitive detection methods (Friedman et al., 2009). Furthermore, recent data show that miRNAs can also act through binding in the ORF of coding genes (Baek et al., 2008; Selbach et al., 2008; Tay et al., 2008), thereby further increasing the degree of miRNA control of the transcriptome.

Advances in quantitative mass spectrometry have enabled the study of the impact of miRNAs on the proteome as a whole, rather than selected targets. Using mass spectrometry, two studies compared the impact of exogenously added miRNAs on the proteome of HeLa cell nuclei (Baek et al., 2008; Selbach et al., 2008). Additionally, these groups analyzed the impact of removing a miRNA, either by deletion in the mouse (Baek et al., 2008) or by endogenous miRNA knockdown in HeLa cells (Selbach et al., 2008). These complimentary approaches revealed several important findings: First, bioinformatic analysis of downregulated genes confirmed that the most important predictor of target recognition is a sequence match to the miRNA seed sequence in the 3' UTR. Second, translational repression is more often than not associated with a parallel decrease in mRNA levels. Third, miRNA effects on their targets are rarely dramatic, instead resulting in small changes in levels across many proteins (Baek et al., 2008; Selbach et al., 2008).

A surprising additional role for microRNAs activating, rather than suppressing, translation has been reported. Specifically, when cells in culture were induced to exit the cell cycle by serum starvation, miR-369-3 and let-7a bound the 3' UTRs and upregulated translation of TNF α and HMGA2 (Vasudevan et al., 2007). Under cell growth conditions, these same miRNAs suppressed the targets suggesting a switch between miRNA induced suppression and activation of translation depending on the cell cycle status (Vasudevan et al., 2007). Transcriptional activation has also been proposed as an alternative role for miRNAs. Following introduction of exogenously synthesized dsRNAs directed to their promoters, expression of E-cadherin, p21, VEGF, and the progesterone receptor were increased (Janowski et al., 2007; Li et al., 2006). The mechanism for such upregulation may involve the dsRNAs targeting antisense transcripts produced at these promoters and the recruitment of Ago1 (Schwartz et al., 2008). An endogenous miRNA, miR-373, may act in a similar fashion to these exogenously introduced dsRNAs suggesting a potential physiological role for such a mechanism (Place et al., 2008). It will be interesting to see how robust these novel mechanisms of miRNA action are in vivo.

3. The role of small RNAs in embryonic stem cells

The first miRNA, lin-4, was identified based on the alterations in cell lineage specification of *C. elegans* (Lee et al., 1993). Analysis of many other organisms has confirmed a central role of miRNAs in normal metazoan development. In mammals, multiple components of the miRNA pathway are essential and their deletion results in early

embryonic lethality. This embryonic lethality has been observed for the deletion of *Dicer* (Bernstein et al., 2003), *Dgcr8* (Wang et al., 2007), and *Ago2* (Alisch et al., 2007; Liu et al., 2004; Morita et al., 2007). To gain insight into early development, embryonic stem cells that contained null alleles of *Dgcr8* and *Dicer* were generated (Kanellopoulou et al., 2005; Murchison et al., 2005; Wang et al., 2007). ES cells lacking either *Dgcr8* or *Dicer* have defects in proliferation and differentiation. The *Dicer* Δ/Δ ES cells appeared to have more severe phenotypes than the *Dgcr8* Δ/Δ ES cells, including a stricter block in differentiation, a severer proliferation defect immediately following loop out, and an inability to form teratomas. These differences suggested the presence of *Dgcr8*-independent, *Dicer*-dependent small RNAs – a conclusion that has been recently confirmed (Babiarz et al., 2008).

Although these phenotypes occurred in the absence of all canonical miRNAs, it remained possible that they could be explained at least in part by one or a few miRNAs that are highly expressed in ES cells. For example, one important difference between wild type and *Dgcr8* Δ/Δ ES cells is that the ES cells accumulated in the G1 phase of the cell cycle in mutants (Wang et al., 2007). Initial attempts at uncovering miRNAs responsible for this defect in a wild-type background failed, presumably due to redundancy among the ES cell miRNAs. Supporting this notion, relatively few individual miRNA knockouts in *C. elegans* have defects in development or viability (Miska et al., 2007). To circumvent the issue of redundancy an alternative approach was taken. Starting with the *Dgcr8* Δ/Δ ES cells lacking all canonical miRNAs, individual miRNAs were added back in the form of miRNA mimics, chemically synthesized double stranded miRNAs representing the post-*Dicer* miRNA duplex. This approach enabled the study of each miRNA in isolation from the normal milieu. A screen using miRNA mimics representing 266 of the 461 annotated miRNAs (Griffiths-Jones et al., 2008) revealed a large family of miRNAs, sharing the same sequence that could rescue the G1 defect in *Dgcr8* Δ/Δ ES cells (Wang et al., 2008). This family includes a subset of the *mir-290* cluster of miRNAs, which are specifically expressed in ES cells (Houbaviy et al., 2003; Wang et al., 2008). These miRNAs were coined ESCC for ES cell-specific cell cycle-regulating miRNAs. Further analysis uncovered targets for the ESCC miRNAs that included multiple inhibitors of the G1/S transition, regulators of the cyclin/cdk2 pathway, including p21 (Cdkn1a), Rb, Rbl2, and Lats2 (Wang et al., 2008). These results correlated nicely with previous results with p21, which showed that while its mRNA levels changed little, its protein levels rapidly increased upon differentiation coincidentally with the lengthening of the G1 phase of the cell cycle (Sabapathy et al., 1997). Taken together, these findings show that in ES cells, the *mir-290* cluster is highly expressed, members of which suppress inhibitors of the G1/S transition promoting an abbreviated G1 phase and hence a shortened cell cycle. Upon differentiation, the *mir-290* cluster is rapidly down regulated, allowing the translation of the G1/S inhibitors and the lengthening of G1.

In addition to the role the *mir-290* cluster plays in the unique ES cell cycle, two groups identified an indirect control of DNA methyltransferase by the *mir-290* cluster (Sinkkonen et al., 2008). Similar to Wang et al, the authors found that the *mir-290* cluster controlled Rbl2. The authors also found that members of the DNA methyltransferase complex, Dnmt3a and 3b, were downregulated in *Dicer* Δ/Δ relative to wild type. Rbl2 had previously been shown to transcriptionally suppress *Dnmt3b* (Litovchick et al., 2007) and the authors showed that inhibition of Rbl2 in the knockout cells rescues Dnmt3a and 3b levels. The authors suggest that the diminished Dnmt levels explain the deficiency of *Dicer* null cells to differentiate as de novo DNA methylation is important in suppressing pluripotency genes (Feldman et al., 2006). However, unlike the *Dgcr8* and *Dicer* knockouts, differentiation appears to initiate normally and pluripotency genes are downregulated in the absence of DNMT3a and 3b, with a fraction of cells reactivating the pluripotency program when returned to ES cell conditions (Feldman et al., 2006). Furthermore, the *mir-290* cluster members do not rescue differentiation (Melton and Blelloch, unpublished observation). In either case, the de novo DNMTs appear to be important indirect targets of the *mir-290* cluster in ES cells.

Taken together, the study of ES cells that lacked miRNAs specifically (*dgcr8* Δ/Δ) or even broader classes of small RNAs (*dicer* Δ/Δ) have been used successfully to uncover important roles for the *mir-290* cluster of miRNAs: ES cell cycle regulation and DNA methylation. Although earlier studies had identified the *mir-290* cluster as highly enriched in ES cells and the early embryo (Houbaviy et al., 2003), the precise function of these abundant miRNAs had not been appreciated. The combination of global miRNA knockout models along with miRNA mimics and inhibitors should continue to identify miRNAs responsible for specific phenotypes in ES cells and other differentiated cell types.

4. Transcriptional regulation of the miRNA content of a cell

Growing evidence suggests that, like the transcriptome and proteome, the miRNA content of an individual cell is unique (Houbaviy et al., 2003; Marson et al., 2008). For instance, ES cells are predominated by the *mir-290*

cluster (Houbaviy et al., 2003; Marson et al., 2008) and a more differentiated cell type, such as neural precursor cells (NPCs), are predominated by the let-7 family (Marson et al., 2008). Several recent studies have begun to shed light into regulation of the miRNA content of a cell (Marson et al., 2008; Ozsolak et al., 2008). Not surprisingly, many of the mechanisms appear similar to the regulation of other developmental genes. Much of the work has focused on ES cells, but the themes of the findings presented thus far are likely operating in other tissue types.

For instance, Marson and colleagues recently published evidence that the core pluripotency transcription factors, Oct4, Sox2, Nanog, and Tcf3, are present at the promoter of the ES-cell specific *mir-290* and *mir-302* clusters in mouse ES cells. Additionally, these authors showed that Oct4 is present at the human ES-cell specific *mir-302* cluster. Intellectually, this result is very satisfying, as it ties the *mir-290* cluster into the ES cell transcriptional network. Indeed, knockdown of *Oct4* results in decreased expression of the *mir-290* cluster. Furthermore, during ES cell differentiation, the down-regulation of the *mir-290* cluster is coincident with the loss of Oct4, Nanog, Sox2, and Tcf3.

Beyond identifying how the core ES-cell transcription factor machinery controls certain miRNAs, Marson and colleagues also identified mechanisms for how certain miRNAs are epigenetically silenced in ES cells. Previous work in mouse ES cells had identified a surprising epigenetic domain, coined a bivalent domain, which contained both active (Histone H3K4me3) and repressive (Histone H3K27me3) marks in the same genomic region (Bernstein et al., 2006). These bivalent domains were predominantly found at the promoters of developmentally important transcription factors that were silent in ES cells, but became activated with differentiation in a lineage dependent fashion (Bernstein et al., 2006). The bivalent domains appear to “resolve” with differentiation: genes that are activated retain the H3K4me3 and lose the H3K27me3 mark, while genes that are suppressed lose H3K4me3 and retain H3K27me3 (Bernstein et al., 2006; Mikkelsen et al., 2007). By examining the H3K4me3 and H3K27me3 marks at miRNA promoters in ES cells and more differentiated cell types, Marson and colleagues were able to show that these bivalent domains also occur at miRNA genes that are silent in ES cells and are activated in a lineage specific fashion (Marson et al., 2008). The most compelling example the authors provide is that of *mir-9*, which is present in NPCs but absent in ES cells and embryonic fibroblasts (MEFs). The promoter of *mir-9* is bivalent in ES cells and is resolved to H3K4me3 in NPCs, versus H3K27me3 in MEFs.

Taken together, these data suggest that regulation of miRNAs during development follows many of the same rules as other developmentally important genes. The unique content of transcription factors and epigenetic modifications found in any particular cell will define the diversity of miRNAs present. In turn, these miRNAs feed back to regulate the signaling pathways, transcription factors and epigenetic status of genes resulting in both negative and positive feedback loops (Marson et al., 2008). Therefore, during cell fate transitions these programs will change in parallel reinforcing the shift from one cell type to another.

5. Post-transcriptional regulation of the processing of miRNAs

Beyond transcription, there are additional layers of regulation of miRNA content of cells. In particular, there is growing evidence for regulation occurring at the level of miRNA biogenesis (Table 1). This regulation, could occur at any one of the steps in their biogenesis pathway including Microprocessor cleavage, Exportin-5 nuclear/cytoplasmic export, Dicer cleavage, transfer to RISC, and RISC silencing. At the global level, it is now clear that the levels of the Microprocessor are very tightly regulated. This regulation occurs through a combination of a negative and a positive feedback loop (Han et al., 2009). The negative feedback loop consists of the Microprocessor cleaving hairpins in the 5'UTR and coding sequence of *Dgcr8*, thereby destabilizing its mRNA. The positive feedback loop involves *Dgcr8* protein stabilizing the Drosha protein through an unclear mechanism. The end result is that any transient shifts in the Microprocessor levels will be rapidly corrected. However, Microprocessor levels can differ between cells both during normal development (Shiohama et al., 2003) and in disease (Ambs et al., 2008; Merritt et al., 2008; Muralidhar et al., 2007), suggesting a resetting of the homeostatic Microprocessor levels.

At the individual miRNA level, there is mounting evidence for both positive and negative regulation of biogenesis. For example, analysis of TGF- β and BMP4 activation of primary human pulmonary artery smooth muscle cells revealed that miR-21 levels are increased post-transcriptionally following growth factor addition (Davis et al., 2008). Because the SMAD proteins are direct targets of TGF- β and BMP4, the authors initially investigated whether receptor-specific SMADs would induce transcription of the pri-miR-21 transcript. However, the pri-miR-21 levels remained unchanged while the pre-, and mature miR-21 levels rose following TGF- β or BMP4 treatment. To confirm a lack of transcription was modulating miRNA levels, the cells were treated with both BMP4 and α -amanitin, which inhibits RNA polymerase II. In the presence of α -amanitin, there was a similar rise in pre-miR-21 and mature miR-21 levels, but an absence

Table 1. miRNAs exhibiting post-transcriptionally regulated processing

miRNA	Biogenesis Step	Type of Regulation	Factor	Cell Type	References
miR-21	Microprocessor	Promotes	SMADs	primary human pulmonary artery smooth muscle cells	(Davis et al., 2008)
miR-18a	Microprocessor	Promotes	hnRNP A1		(Guil and Caceres 2007; Michlewski et al. 2008)
let-7 family	Microprocessor	Inhibits	Lin-28	mES and mEC	(Newman et al. 2008; Viswanathan et al. 2008)
let-7 family	Dicer	Inhibits	Lin-28	mES	(Rybak et al. 2008)
let-7 family	Dicer	Inhibits	Lin-28	Hepatocellular	(Heo et al. 2008)
			Lin-28B	Carcinoma Cell Lines	
miR-221, miR-222	RISC translational Inhibition of p27	Inhibits	Dnd1	Human Germ Cells	(Kedde et al. 2007)
miR-138	Dicer	Inhibits	Unknown	HeLa	(Obenosterer et al. 2006)
miR-101-1, let-7a-1, miR-379, miR-31	Microprocessor	Promotes	Unknown	In vitro	(Michlewski et al. 2008)

of transcription of a known downstream BMP4 target. SMAD1 was previously shown to interact with p68 (Warner et al., 2004), an RNA-helicase present in the Microprocessor complex. Furthermore, it was shown that the loss of p68 causes a decrease in miR-21 in E9.5 embryos (Fukuda et al., 2007). Therefore, Davis et al. tested and showed that the SMADs interact with pri-mir-21, recruit the Microprocessor via the p68 subunit, thereby enhance pri- to pre-mir-21 processing (Davis et al., 2008). These results showed that signaling molecules are able to direct post-transcriptional increases in miRNA processing, which enables rapid miRNA mediated responses.

Another example of positive post-transcriptional regulation of miRNAs is the ribonuclear protein, hnRNP A1, which regulates the processing of miR-18a. Using CLIP (cross-linking and IP) (Ule et al., 2005; Ule et al., 2003) to identify targets of hnRNP A1, Guil et al. pulled down miR-18a and showed that binding of hnRNP A1 to the pri-miR-18a transcript facilitates its processing (Guil and Caceres, 2007). Depletion of hnRNP A1 in HeLa cells resulted in a reduction in the amounts of miR-18a (Guil and Caceres, 2007). A follow-up study showed that hnRNP A1 binds to the stem of pri-miR-18a, altering the secondary structure of the hairpin, which facilitates cleavage by the Microprocessor (Michlewski et al., 2008). The authors suggest this type of regulation may be widespread, as they found evolutionary conservation in the stem loops of a variety of pre-miRNA hairpins suggesting selection to keep the loop intact, and hence a likely regulatory role. Furthermore, anti-sense oligonucleotides against the conserved loop of the pre-miRNA hairpins often diminished processing in vitro (Michlewski et al., 2008). These results suggest that there may be an entire class of proteins that bind the loop of the pri-miRNA hairpins in order to promote Microprocessor cleavage. The lack of a Microprocessor facilitating protein may allow pri-miRNAs to accumulate and then be rapidly processed to mature miRNAs following the expression of that facilitator.

In contrast to the hnRNP story, another protein, Lin-28, has been shown to block miRNA processing. The exact step where the processing is blocked has been controversial as two studies identified a block at the Microprocessor step (Newman et al., 2008; Viswanathan et al., 2008) while two others identified the block at the Dicer step (Heo et al., 2008; Rybak et al., 2008). Importantly though, they all agree that Lin-28 inhibits the maturation of a particular subset of miRNAs, the let-7 miRNAs. The search for Lin-28 began with the finding that while mature let-7 was absent in ES cells and the early embryo, the pri-miRNA was being transcribed (Thomson et al., 2006; Wulczyn et al., 2007). With ES cell differentiation or embryonic maturation, mature let-7 levels rose with little change in the steady state pri-let-7 levels. This finding suggested a block in processing somewhere between the pri- and mature miRNA. As the pri-let-7 miRNAs were present in the ES cells, two groups hypothesized that the block had to be at the pri- to pre-miRNA processing step (Newman et al., 2008; Viswanathan et al., 2008). They determined that there was indeed a protein present in ES, but not differentiated cell extracts that could block pri- to pre-miRNA processing. Therefore, they used affinity columns along with an in vitro Microprocessor activity assay to identify the responsible protein(s). The affinity columns consisted of beads linked to either pre-let-7g (Viswanathan et al., 2008) or the let-7d loop (Newman

et al., 2008). Purified extracts from these columns retained inhibitory activity and, therefore, were analyzed by mass spectrometry. From this analysis and follow-up experiments, both groups identified Lin-28, and the closely related Lin-28B protein as the inhibitory factors. Indeed, ectopic expression of Lin-28 in a cell line that has high mature let-7 levels and low Lin-28 resulted in a decrease in mature let-7, while Lin-28 knockdown in ES cells resulted in elevated mature let-7 levels (Newman et al., 2008; Viswanathan et al., 2008). These findings led the authors to conclude that Lin-28 bound to the *let-7* hairpins in the pri-miRNA preventing Microprocessor cleavage.

However, Lin-28 is predominantly a cytoplasmic protein (Balzer and Moss, 2007), while pri- to pre- processing occurs in the nucleus. Work by two other groups may have shed some light on this conundrum. Rybak and colleagues had previously suggested that pre-, not pri-let-7 family members accumulated in ES cells (Wulczyn et al., 2007). In vitro assays using the pre-let-7 hairpin as opposed to the longer pri-let-7, revealed that Lin-28 blocks mature let-7 accumulation at the Dicer step (Rybak et al., 2008). Additionally, Heo et al. found that Lin-28B blocks Dicer cleavage of pre-let-7 by promoting poly-uridylation of the pre-let-7. This uridylation blocks Dicer cleavage and simultaneously targets the RNA for degradation. Further, they found that the let-7 inhibitory function was present primarily in the cytoplasm, consistent with Lin-28's subcellular localization.

Recent evidence has emerged that suggests Lin-28, and by extension, let-7 family members, are integrated into the pluripotency circuit. The *Lin-28* promoter is occupied by the core pluripotency transcription factors Oct4, Nanog, Sox2, and Tcf3 (Marson et al., 2008). The high level of Lin-28 expression prevents the accumulation of mature let-7 family miRNAs. Surprisingly, the promoter of at least one member of the let-7 family, let-7g, is also bound by Oct4, Nanog, Sox2, and Tcf3, likely driving high levels of transcription of pri-let-7g in ES cells (Marson et al., 2008). Upon the induction of differentiation, Oct4, Nanog, and Sox2 are downregulated. This results in the downregulation of Lin-28 and concomitant upregulation of the processing of the accumulated pri- or pre-let-7g to mature let-7g. This upregulation of let-7 further downregulates Lin-28 by targeting the Lin-28 3' UTR (John et al., 2004; Moss and Tang, 2003; Nelson et al., 2004; Reinhart et al., 2000; Wu and Belasco, 2005) producing a positive feedback loop or “bistable switch” (Heo et al., 2008). This type of regulatory circuit allows the powerful and rapid transition from a pluripotent stem cell that lacks let-7 to a more differentiated cell that has mature let-7.

Finally, a recent report revealed that following processing and RISC loading, at least one RNA binding protein is capable of inhibiting the silencing effect of RISC on certain mRNAs. Noting the a high level of conservation of the 3' UTR of p27, outside of the miR-221 and miR-222 seeds, Kedde and colleagues speculated that an RNA-binding protein may recognize this sequence and modulate miR-221/222 silencing of p27 (Kedde et al., 2007). Through co-transfections of candidate RNA-binding proteins and miR-221 in HEK-293 cells, the authors found Dead end 1 (Dnd1) prevented miR-221 silencing of p27. Surprisingly, mutating the miR-221 site to a perfect match, showed that Dnd1 could not prevent the siRNA slicing of the transcript. This finding suggests that Dnd1 does not merely function through steric hindrance, although it may prevent a stable, long-term association of RISC with the 3' UTR. The authors conclusively showed that Dnd1 inhibits miR-221 silencing of p27. It will be interesting to see if Dnd1 or similar proteins influence other mRNA/miRNA interactions.

Taken together, the findings discussed above show a broad range of control of miRNAs at nearly every step in their biogenesis as well as execution of their function. At this point, descriptions of only a handful miRNAs controlled at any particular step are published. It will be fascinating to see how many miRNAs are controlled at each of these steps, and how many proteins are capable of these types of regulation.

6. miRNAs and induced pluripotency

The finding that a combination of transcription factors can reprogram differentiated adult cells into a pluripotent state has revolutionized the regenerative medicine field. Interestingly, alternative combinations of viral expressed factors have been used to successfully reprogram mouse and human ES cells (Takahashi et al., 2007; Takahashi and Yamanaka, 2006; Yu et al., 2007). The original mix of factors used to dedifferentiate mouse somatic cells to induced pluripotent stem cells was a combination of Oct4, Sox2, c-Myc, and Klf4 (Takahashi and Yamanaka, 2006). This same set of factors was later shown to produce human iPS cells as well (Lowry et al., 2008; Park et al., 2008; Takahashi et al., 2007). However, another group identified a different combination of factors that could reprogram human cells consisting of Oct4, Sox2, Nanog, and Lin-28 (Yu et al., 2007). From a “miRNA-centric” view of reprogramming, these reprogramming factors are extremely intriguing. Oct4 and Sox2 (and Nanog) are important for driving the ES cell-specific miRNA clusters *mir-290* (mouse) and *mir-302* (mouse and human) (Marson et al., 2008). Lin-28 inhibits let-7 biogenesis, and, therefore, may support reprogramming by depleting the cells of mature




Figure 2. The Non-canonical miRNA biogenesis pathway. Mirtrons are short introns which form pre-miRNA hairpins following splicing and debranching of the transcript. The endogenous shRNAs are directly transcribed as pre-miRNA hairpins. Adapted from Babiarz et al., 2008.

let-7. Finally, c-Myc controls the transcription of many miRNAs, in both a positive and negative fashion. c-Myc can drive the expression of the *mir-17-92* cluster (O'Donnell et al., 2005), which is one of the more highly expressed families in ES cells (Calabrese et al., 2007; Marson et al., 2008). Additionally, c-Myc can downregulate a number of miRNAs, including let-7 family members (Chang et al., 2008). Therefore, the reprogramming factors may be acting in part by altering the miRNA content of the somatic cell to that of a pluripotent stem cell in order to induce dedifferentiation.

Indeed, a recent report has shown that the transfection of a subset of the *mir-290* cluster miRNAs can replace Myc in reprogramming mouse fibroblasts to induced pluripotent stem cells (Judson et al., 2009). These were the same ESCC miRNAs that were previously shown to promote the ES cell cycle (Wang et al., 2008). This finding suggests that the transcriptional activation of the *mir-290* cluster and possibly the establishment of the unique ES cell cycle structure are critical steps in de-differentiation. Indeed, both c-Myc and n-Myc bind the promoter of the *mir-290* cluster, suggesting that one of the critical roles of Myc in reprogramming is to activate these miRNAs. However, there must be an intermediate step since the promoter of the *mir-290* cluster is epigenetically silenced in fibroblasts (H3K27me3 positive, H3K4me3 negative). Importantly, the ESCC miRNAs do not recapitulate all the downstream affects of Myc. In particular, unlike Myc, the ESCC miRNAs do not promote proliferation of MEFs and they produce a homogenous population of fully reprogrammed iPS colonies. Furthermore, teratomas from Myc-iPS, but not the ESCC miRNA-iPS cells, had a tendency to be invasive. Together, these findings show how microRNAs can play a central role in induced pluripotency.

7. Non-canonical miRNAs and endogenous siRNAs

While most miRNAs require both the Microprocessor and Dicer for their biogenesis, exceptions have been recently uncovered (Figure 2). For example, deep sequencing of small RNAs in *Drosophila* uncovered a novel subclass of miRNAs, the mirtrons (Okamura et al., 2007; Ruby et al., 2007). The mirtrons are generated from short introns that fold into pre-miRNA hairpins following debranching of the intron lariat. These mirtrons were discovered soon after in mammals (Babiarz et al., 2008; Berezikov et al., 2007). Babiarz and colleagues discovered the mirtrons in mouse ES cells by analyzing all small RNAs whose biogenesis relies on Dicer, but not the Microprocessor (Babiarz et al., 2008). However, the mirtrons made up a small fraction of small RNAs and as such, their specific role remains unclear. Most




Figure 3. The endo-siRNA biogenesis pathway. Hairpin siRNAs are transcribed and fold into a long hairpin, which is cleaved multiple times by Dicer. cis-siRNAs are derived from transcription on both strands of DNA, resulting in a two-strand dsRNA with perfect complementarity. trans-siRNAs are derived from gene/pseudogene pairs from distant sites in the genome. Because of mutations in the pseudogene, complementarity of the dsRNA is not perfect. hp-siRNAs have been found in mouse oocytes and ES cells. trans-siRNAs and cis-siRNAs have only been found in oocytes. Adapted from Babiarz et al., 2008.

of the Dicer-dependent, Dgcr8-independent small RNAs were derived from intergenic regions of the genome. The two most abundant of these miRNAs, miR-320 and miR-484 had been previously annotated and thought to be canonical miRNAs. Careful analysis of these two loci, as well as comparison to another related locus, suggest that miR-320 and miR-484 may be transcribed directly as short hairpin RNAs, similar to exogenously introduced short hairpins used experimentally by many labs (Dykxhoorn and Lieberman, 2005; Elbashir et al., 2001; Hannon and Rossi, 2004). However, it cannot be ruled out that there is a yet unknown nuclease(s) that act on the transcripts from these loci to produce the pre-miRNA. The Dgcr8-independence of these miRNAs was recently confirmed in mouse embryonic skin (Yi et al., 2009). These findings make it clear that there is a diversity of biogenic mechanisms in place in metazoan cells that can produce miRNAs. Whether these miRNAs act in the same manner as the canonical miRNAs is an open question.

Interestingly, both *miR-320* and *miR-484* are found upstream of genes and expressed in the antisense direction to those genes. Therefore, it was hypothesized they may be regulating the promoter regions of their respective genes. Indeed, in HeLa cells, where miR-320 levels are relatively high, H3K27me3 was found in the promoter of the adjacent gene. Additionally, the exogenous introduction of miR-320 into HEK-293 cells, which normally have undetectable miR-320, enhances H3K27me3 within the promoter of its adjacent gene, suppressing expression (Kim et al., 2008). However, analysis of genome-wide H3K27me3 in mouse ES cells, embryonic fibroblasts, and neural precursor cells (Mikkelsen et al., 2007) revealed no enrichment of H3K27me3 at this locus even though miR-320 is clearly present (Babiarz and Blelloch, unpublished observation). Therefore, it is unclear whether this role of miR-320 may be cell context dependent.

In addition to identifying Microprocessor-independent miRNAs, the deep sequencing of small RNAs in the different mutant background uncovered endogenous siRNAs (endo-siRNAs) in mES cells (Babiarz et al., 2008) (Figure 3). The presence of endo-siRNAs in mammalian cells was thought unlikely, because the long, double stranded precursors to endo-siRNAs would trigger the interferon response to dsRNAs (Stark et al., 1998). However, there are at least two known cell types where the interferon response does not occur: oocytes and ES cells (Wianny and Zernicka-Goetz, 2000; Yang et al., 2001). Indeed, along with the Babiarz et al. study in ES cells, two other reports identified mammalian endo-siRNAs, but in mouse oocytes (Tam et al., 2008; Watanabe et al., 2008). In the mouse oocytes, endo-siRNAs arose from two primary sources: retrotransposons, both LINE L1 and LTR elements, and

gene/pseudogene pairs. A loss of Dicer in oocytes results in an increase in retrotransposons that have endo-siRNAs associated with them and an increase in transcripts associated with the gene/pseudogene pairs (Tam et al., 2008; Watanabe et al., 2008). However, it is currently unclear if the increase in transcripts is due to a loss of Dicer degrading the dsRNAs, or if it is a secondary effect of siRNAs normally feeding back to destabilize their transcripts of origin.

In the mES cells, a very different population of endo-siRNAs was uncovered (Babiarz et al., 2008). The ES cells did not contain any endo-siRNAs arising from gene/pseudogene pairs; instead, the endo-siRNAs arose from SINE B1/Alu elements, most of which could be mapped back to two loci within the genome (Babiarz et al., 2008). These loci contained two tandem, inverted SINE B1/Alu elements, that when transcribed form a long hairpin. Dicer could then processively cleave this hairpin with an approximate 21 nucleotide (nt) periodicity, giving rise to a number of unique small RNAs. The role of these endo-siRNAs is unclear; the most straightforward mechanism of endo-siRNA function, degradation of SINE RNAs, does not appear to occur in ES cells. The loss of Dicer in mES cells does not result in an increase in SINE transcripts (Calabrese et al., 2007).

Mammalian endo-siRNAs are developmentally regulated. As described above, there was no overlap between the species of endo-siRNAs found in oocytes and ES cells (Babiarz et al., 2008). Whether the levels of the different endo-siRNAs are regulated transcriptionally and/or post-transcriptionally is currently unknown. It is possible that demethylation during germ cell development enable expression of the pseudogenes resulting in siRNA production which then play a critical role in the oocyte to suppress the function of the pseudogene's partner. Then, as the pseudogenes are methylated and silenced during early embryogenesis, the gene partners are stabilized and translated. Similarly, the presence of LINE and LTR siRNAs in oocytes but not ES cells may represent patterns in their expression. Interestingly, LINE L1 and IAP mRNAs are increased in *Dicer* Δ/Δ mES cells (Kanellopoulou et al., 2005), even though no Dicer-dependent small RNAs are generated from those retrotransposons (Babiarz et al., 2008). How are these retrotransposons controlled? Can the SINE endo-siRNAs control the LINE L1 and IAP elements? Or, are there other, indirect methods, such as a decrease in DNA methylation due to the down-regulation of Dnmt3a/b through Rbl2 (Benetti et al., 2008; Sinkkonen et al., 2008)? Furthermore, it remains unknown whether there are endo-siRNAs in somatic tissues and, if so, whether they are unique from those uncovered in oocytes and ES cells suggesting regulatory roles for endo-siRNAs throughout development.

8. Conclusions

The study of small RNAs has been occurring at a fast pace with many exciting discoveries being uncovered. The importance of miRNAs in development is clear, with severe phenotypes associated with their loss (Bernstein et al., 2003; Kanellopoulou et al., 2005; Wang et al., 2007). However, the mechanism of action of miRNAs in mammalian development remains largely unknown and hotly pursued. Although a great deal has been learned about how miRNAs are generated, many questions remain unanswered. Chief among those is what are the precise roles of individual miRNAs in any particular cell. How are miRNA levels controlled? Are miRNAs in the same family redundant, or are there non-overlapping functions of the different family members. How central is the miRNA content of a cell in determining the cell's identity? What is the developmental role of endogenous siRNAs? Tackling these and other questions will keep the miRNA field a productive area of study for years to come.

9. References

- Alisch, R. S., Jin, P., Epstein, M., Caspary, T., and Warren, S. T. (2007). Argonaute2 is essential for mammalian gastrulation and proper mesoderm formation. *PLoS Genet* 3, e227.
- Ambs, S., Prueitt, R. L., Yi, M., Hudson, R. S., Howe, T. M., Petrocca, F., Wallace, T. A., Liu, C. G., Volinia, S., and Calin, G. A., et al. (2008). Genomic profiling of microRNA and messenger RNA reveals deregulated microRNA expression in prostate cancer. *Cancer Res* 68, 6162–6170.
- Babiarz, J. E., Ruby, J. G., Wang, Y., Bartel, D. P., and Blelloch, R. (2008). Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes Dev* 22, 2773–2785.
- Baek, D., Villen, J., Shin, C., Camargo, F. D., Gygi, S. P., and Bartel, D. P. (2008). The impact of microRNAs on protein output. *Nature* 455, 64–71.
- Balzer, E., and Moss, E. G. (2007). Localization of the developmental timing regulator Lin28 to mRNP complexes, P-bodies and stress granules. *RNA Biol* 4, 16–25.

- Bartel, D. P. (2009). MicroRNAs: Target Recognition and Regulatory Functions. *Cell* 136, 215–233.
- Benetti, R., Gonzalo, S., Jaco, I., Munoz, P., Gonzalez, S., Schoeftner, S., Murchison, E., Andl, T., Chen, T., and Klatt, P., et al. (2008). A mammalian microRNA cluster controls DNA methylation and telomere recombination via Rbl2-dependent regulation of DNA methyltransferases. *Nat Struct Mol Biol* 15, 268–279.
- Berezikov, E., Chung, W. J., Willis, J., Cuppen, E., and Lai, E. C. (2007). Mammalian mirtron genes. *Mol Cell* 28, 328–336.
- Bernstein, B. E., Mikkelsen, T. S., Xie, X., Kamal, M., Huebert, D. J., Cuff, J., Fry, B., Meissner, A., Wernig, M., and Plath, K., et al. (2006). A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* 125, 315–326.
- Bernstein, E., Kim, S. Y., Carmell, M. A., Murchison, E. P., Alcorn, H., Li, M. Z., Mills, A. A., Elledge, S. J., Anderson, K. V., and Hannon, G. J. (2003). Dicer is essential for mouse development. *Nat Genet* 35, 215–217.
- Bohnsack, M. T., Czapinski, K., and Gorlich, D. (2004). Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA* 10, 185–191.
- Cai, X., Hagedorn, C. H., and Cullen, B. R. (2004). Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* 10, 1957–1966.
- Calabrese, J. M., Seila, A. C., Yeo, G. W., and Sharp, P. A. (2007). RNA sequence analysis defines Dicer's role in mouse embryonic stem cells. *Proc Natl Acad Sci U S A* 104, 18097–18102.
- Carmell, M. A., Xuan, Z., Zhang, M. Q., and Hannon, G. J. (2002). The Argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes Dev* 16, 2733–2742.
- Chang, T. C., Yu, D., Lee, Y. S., Wentzel, E. A., Arking, D. E., West, K. M., Dang, C. V., Thomas-Tikhonenko, A., and Mendell, J. T. (2008). Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat Genet* 40, 43–50.
- Chendrimada, T. P., Gregory, R. I., Kumaraswamy, E., Norman, J., Cooch, N., Nishikura, K., and Shiekhattar, R. (2005). TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature* 436, 740–744.
- Davis, B. N., Hilyard, A. C., Lagna, G., and Hata, A. (2008). SMAD proteins control DROSHA-mediated microRNA maturation. *Nature* 454, 56–61.
- Denli, A. M., Tops, B. B., Plasterk, R. H., Ketting, R. F., and Hannon, G. J. (2004). Processing of primary microRNAs by the Microprocessor complex. *Nature* 432, 231–235.
- Dykxhoorn, D. M., and Lieberman, J. (2005). The silent revolution: RNA interference as basic biology, research tool, and therapeutic. *Annu Rev Med* 56, 401–423.
- Elbashir, S. M., Harborth, J., Lendeckel, W., Yalcin, A., Weber, K., and Tuschl, T. (2001). Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 411, 494–498.
- Feldman, N., Gerson, A., Fang, J., Li, E., Zhang, Y., Shinkai, Y., Cedar, H., and Bergman, Y. (2006). G9a-mediated irreversible epigenetic inactivation of Oct-3/4 during early embryogenesis. *Nat Cell Biol* 8, 188–194.
- Friedman, R. C., Farh, K. K., Burge, C. B., and Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 19, 92–105.
- Fukuda, T., Yamagata, K., Fujiyama, S., Matsumoto, T., Koshida, I., Yoshimura, K., Mihara, M., Naitou, M., Endoh, H., and Nakamura, T., et al. (2007). DEAD-box RNA helicase subunits of the Drosha complex are required for processing of rRNA and a subset of microRNAs. *Nat Cell Biol* 9, 604–611.
- Gregory, R. I., Chendrimada, T. P., Cooch, N., and Shiekhattar, R. (2005). Human RISC couples microRNA biogenesis and posttranscriptional gene silencing. *Cell* 123, 631–640.

- Gregory, R. I., Yan, K. P., Amuthan, G., Chendrimada, T., Doratotaj, B., Cooch, N., and Shiekhattar, R. (2004). The Microprocessor complex mediates the genesis of microRNAs. *Nature* **432**, 235–240.
- Griffiths-Jones, S., Saini, H. K., van Dongen, S., and Enright, A. J. (2008). miRBase: tools for microRNA genomics. *Nucleic Acids Res* **36**, D154–158.
- Grimson, A., Farh, K. K., Johnston, W. K., Garrett-Engele, P., Lim, L. P., and Bartel, D. P. (2007). MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol Cell* **27**, 91–105.
- Grimson, A., Srivastava, M., Fahey, B., Woodcroft, B. J., Chiang, H. R., King, N., Degnan, B. M., Rokhsar, D. S., and Bartel, D. P. (2008). Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature* **455**, 1193–1197.
- Guil, S., and Caceres, J. F. (2007). The multifunctional RNA-binding protein hnRNP A1 is required for processing of miR-18a. *Nat Struct Mol Biol* **14**, 591–596.
- Haase, A. D., Jaskiewicz, L., Zhang, H., Laine, S., Sack, R., Gatignol, A., and Filipowicz, W. (2005). TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA silencing. *EMBO Rep* **6**, 961–967.
- Hammond, S. M. (2005). Dicing and slicing: the core machinery of the RNA interference pathway. *FEBS Lett* **579**, 5822–5829.
- Han, J., Lee, Y., Yeom, K. H., Kim, Y. K., Jin, H., and Kim, V. N. (2004). The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* **18**, 3016–3027.
- Han, J., Lee, Y., Yeom, K. H., Nam, J. W., Heo, I., Rhee, J. K., Sohn, S. Y., Cho, Y., Zhang, B. T., and Kim, V. N. (2006). Molecular basis for the recognition of primary microRNAs by the Drosha-DGCR8 complex. *Cell* **125**, 887–901.
- Han, J., Pedersen, J. S., Kwon, S. C., Belair, C. D., Kim, Y. K., Yeom, K. H., Yang, W. Y., Haussler, D., Blelloch, R., and Kim, V. N. (2009). Posttranscriptional crossregulation between Drosha and DGCR8. *Cell* **136**, 75–84.
- Hannon, G. J., and Rossi, J. J. (2004). Unlocking the potential of the human genome with RNA interference. *Nature* **431**, 371–378.
- Heo, I., Joo, C., Cho, J., Ha, M., Han, J., and Kim, V. N. (2008). Lin28 mediates the terminal uridylation of let-7 precursor MicroRNA. *Mol Cell* **32**, 276–284.
- Houbaviy, H. B., Murray, M. F., and Sharp, P. A. (2003). Embryonic stem cell-specific MicroRNAs. *Dev Cell* **5**, 351–358.
- Janowski, B. A., Younger, S. T., Hardy, D. B., Ram, R., Huffman, K. E., and Corey, D. R. (2007). Activating gene expression in mammalian cells with promoter-targeted duplex RNAs. *Nat Chem Biol* **3**, 166–173.
- John, B., Enright, A. J., Aravin, A., Tuschl, T., Sander, C., and Marks, D. S. (2004). Human MicroRNA targets. *PLoS Biol* **2**, e363.
- Judson, R. L., Babiarz, J. E., Venere, M., and Blelloch, R. (2009). Embryonic stem cell-specific microRNAs promote induced pluripotency. *Nat Biotechnol*.
- Kanellopoulou, C., Muljo, S. A., Kung, A. L., Ganesan, S., Drapkin, R., Jenuwein, T., Livingston, D. M., and Rajewsky, K. (2005). Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev* **19**, 489–501.
- Kedde, M., Strasser, M. J., Boldajipour, B., Oude Vrielink, J. A., Slanchev, K., le Sage, C., Nagel, R., Voorhoeve, P. M., van Duijse, J., and Orom, U. A., et al. (2007). RNA-binding protein Dnd1 inhibits microRNA access to target mRNA. *Cell* **131**, 1273–1286.

- Khvorova, A., Reynolds, A., and Jayasena, S. D. (2003). Functional siRNAs and miRNAs exhibit strand bias. *Cell* *115*, 209–216.
- Kim, D. H., Saetrom, P., Snove, O. Jr., and Rossi, J. J. (2008). MicroRNA-directed transcriptional gene silencing in mammalian cells. *Proc Natl Acad Sci U S A* *105*, 16230–16235.
- Kim, Y. K., and Kim, V. N. (2007). Processing of intronic microRNAs. *EMBO J* *26*, 775–783.
- Landthaler, M., Yalcin, A., and Tuschl, T. (2004). The human DiGeorge syndrome critical region gene 8 and Its D. melanogaster homolog are required for miRNA biogenesis. *Curr Biol* *14*, 2162–2167.
- Lee, R. C., Feinbaum, R. L., and Ambros, V. (1993). The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* *75*, 843–854.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S., and Kim, V. N. (2003). The nuclear RNase III Drosha initiates microRNA processing. *Nature* *425*, 415–419.
- Lee, Y., Kim, M., Han, J., Yeom, K. H., Lee, S., Baek, S. H., and Kim, V. N. (2004). MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* *23*, 4051–4060.
- Lewis, B. P., Burge, C. B., and Bartel, D. P. (2005). Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* *120*, 15–20.
- Lewis, B. P., Shih, I. H., Jones-Rhoades, M. W., Bartel, D. P., and Burge, C. B. (2003). Prediction of mammalian microRNA targets. *Cell* *115*, 787–798.
- Li, L. C., Okino, S. T., Zhao, H., Pookot, D., Place, R. F., Urakami, S., Enokida, H., and Dahiya, R. (2006). Small dsRNAs induce transcriptional activation in human cells. *Proc Natl Acad Sci U S A* *103*, 17337–17342.
- Litovchick, L., Sadasivam, S., Florens, L., Zhu, X., Swanson, S. K., Velmurugan, S., Chen, R., Washburn, M. P., Liu, X. S., and DeCaprio, J. A. (2007). Evolutionarily conserved multisubunit RBL2/p130 and E2F4 protein complex represses human cell cycle-dependent genes in quiescence. *Mol Cell* *26*, 539–551.
- Liu, J., Carmell, M. A., Rivas, F. V., Marsden, C. G., Thomson, J. M., Song, J. J., Hammond, S. M., Joshua-Tor, L., and Hannon, G. J. (2004). Argonaute2 is the catalytic engine of mammalian RNAi. *Science* *305*, 1437–1441.
- Liu, Y., Taverna, S. D., Muratore, T. L., Shabanowitz, J., Hunt, D. F., and Allis, C. D. (2007). RNAi-dependent H3K27 methylation is required for heterochromatin formation and DNA elimination in *Tetrahymena*. *Genes Dev* *21*, 1530–1545.
- Lowry, W. E., Richter, L., Yachechko, R., Pyle, A. D., Tchieu, J., Sridharan, R., Clark, A. T., and Plath, K. (2008). Generation of human induced pluripotent stem cells from dermal fibroblasts. *Proc Natl Acad Sci U S A* *105*, 2883–2888.
- Lund, E., Guttinger, S., Calado, A., Dahlberg, J. E., and Kutay, U. (2004). Nuclear export of microRNA precursors. *Science* *303*, 95–98.
- Marson, A., Levine, S. S., Cole, M. F., Frampton, G. M., Brambrink, T., Johnstone, S., Guenther, M. G., Johnston, W. K., Wernig, M., and Newman, J., et al. (2008). Connecting microRNA genes to the core transcriptional regulatory circuitry of embryonic stem cells. *Cell* *134*, 521–533.
- Meister, G., Landthaler, M., Patkaniowska, A., Dorsett, Y., Teng, G., and Tuschl, T. (2004). Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol Cell* *15*, 185–197.
- Merritt, W. M., Lin, Y. G., Han, L. Y., Kamat, A. A., Spannuth, W. A., Schmandt, R., Urbauer, D., Pennacchio, L. A., Cheng, J. F., and Nick, A. M., et al. (2008). Dicer, Drosha, and outcomes in patients with ovarian cancer. *N Engl J Med* *359*, 2641–2650.

- Michlewski, G., Guil, S., Semple, C. A., and Caceres, J. F. (2008). Posttranscriptional regulation of miRNAs harboring conserved terminal loops. *Mol Cell* 32, 383–393.
- Mikkelsen, T. S., Ku, M., Jaffe, D. B., Issac, B., Lieberman, E., Giannoukos, G., Alvarez, P., Brockman, W., Kim, T. K., and Koche, R. P., et al. (2007). Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* 448, 553–560.
- Miska, E. A., Alvarez-Saavedra, E., Abbott, A. L., Lau, N. C., Hellman, A. B., McGonagle, S. M., Bartel, D. P., Ambros, V. R., and Horvitz, H. R. (2007). Most *Caenorhabditis elegans* microRNAs are individually not essential for development or viability. *PLoS Genet* 3, e215.
- Morita, S., Horii, T., Kimura, M., Goto, Y., Ochiya, T., and Hatada, I. (2007). One Argonaute family member, Eif2c2 (Ago2), is essential for development and appears not to be involved in DNA methylation. *Genomics* 89, 687–696.
- Morlando, M., Ballarino, M., Gromak, N., Pagano, F., Bozzoni, I., and Proudfoot, N. J. (2008). Primary microRNA transcripts are processed co-transcriptionally. *Nat Struct Mol Biol* 15, 902–909.
- Moss, E. G., and Tang, L. (2003). Conservation of the heterochronic regulator Lin-28, its developmental expression and microRNA complementary sites. *Dev Biol* 258, 432–442.
- Muralidhar, B., Goldstein, L. D., Ng, G., Winder, D. M., Palmer, R. D., Gooding, E. L., Barbosa-Morais, N. L., Mukherjee, G., Thorne, N. P., and Roberts, I., et al. (2007). Global microRNA profiles in cervical squamous cell carcinoma depend on Droscha expression levels. *J Pathol* 212, 368–377.
- Murchison, E. P., Partridge, J. F., Tam, O. H., Cheloufi, S., and Hannon, G. J. (2005). Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad Sci U S A* 102, 12135–12140.
- Nelson, P. T., Hatzigeorgiou, A. G., and Mourelatos, Z. (2004). miRNP:mRNA association in polyribosomes in a human neuronal cell line. *RNA* 10, 387–394.
- Newman, M. A., Thomson, J. M., and Hammond, S. M. (2008). Lin-28 interaction with the Let-7 precursor loop mediates regulated microRNA processing. *RNA* 14, 1539–1549.
- O'Donnell, K. A., Wentzel, E. A., Zeller, K. I., Dang, C. V., and Mendell, J. T. (2005). c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* 435, 839–843.
- Okamura, K., Hagen, J. W., Duan, H., Tyler, D. M., and Lai, E. C. (2007). The mirtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell* 130, 89–100.
- Ozsolak, F., Poling, L. L., Wang, Z., Liu, H., Liu, X. S., Roeder, R. G., Zhang, X., Song, J. S., and Fisher, D. E. (2008). Chromatin structure analyses identify miRNA promoters. *Genes Dev* 22, 3172–3183.
- Park, I. H., Zhao, R., West, J. A., Yabuuchi, A., Huo, H., Ince, T. A., Lerou, P. H., Lensch, M. W., and Daley, G. Q. (2008). Reprogramming of human somatic cells to pluripotency with defined factors. *Nature* 451, 141–146.
- Place, R. F., Li, L. C., Pookot, D., Noonan, E. J., and Dahiya, R. (2008). MicroRNA-373 induces expression of genes with complementary promoter sequences. *Proc Natl Acad Sci U S A* 105, 1608–1613.
- Rand, T. A., Ginalski, K., Grishin, N. V., and Wang, X. (2004). Biochemical identification of Argonaute 2 as the sole protein required for RNA-induced silencing complex activity. *Proc Natl Acad Sci U S A* 101, 14385–14389.
- Rand, T. A., Petersen, S., Du, F., and Wang, X. (2005). Argonaute2 cleaves the anti-guide strand of siRNA during RISC activation. *Cell* 123, 621–629.
- Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., Horvitz, H. R., and Ruvkun, G. (2000). The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403, 901–906.

- Riddle, N. C., and Elgin, S. C. (2008). A role for RNAi in heterochromatin formation in *Drosophila*. *Curr Top Microbiol Immunol* 320, 185–209.
- Ruby, J. G., Jan, C. H., and Bartel, D. P. (2007). Intronic microRNA precursors that bypass Drosha processing. *Nature* 448, 83–86.
- Rybak, A., Fuchs, H., Smirnova, L., Brandt, C., Pohl, E. E., Nitsch, R., and Wulczyn, F. G. (2008). A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* 10, 987–993.
- Sabapathy, K., Klemm, M., Jaenisch, R., and Wagner, E. F. (1997). Regulation of ES cell differentiation by functional and conformational modulation of p53. *EMBO J* 16, 6217–6229.
- Schwartz, J. C., Younger, S. T., Nguyen, N. B., Hardy, D. B., Monia, B. P., Corey, D. R., and Janowski, B. A. (2008). Antisense transcripts are targets for activating small RNAs. *Nat Struct Mol Biol* 15, 842–848.
- Schwarz, D. S., Hutvagner, G., Du, T., Xu, Z., Aronin, N., and Zamore, P. D. (2003). Asymmetry in the assembly of the RNAi enzyme complex. *Cell* 115, 199–208.
- Selbach, M., Schwanhaussner, B., Thierfelder, N., Fang, Z., Khanin, R., and Rajewsky, N. (2008). Widespread changes in protein synthesis induced by microRNAs. *Nature* 455, 58–63.
- Shiohama, A., Sasaki, T., Noda, S., Minoshima, S., and Shimizu, N. (2003). Molecular cloning and expression analysis of a novel gene DGCR8 located in the DiGeorge syndrome chromosomal region. *Biochem Biophys Res Commun* 304, 184–190.
- Sinkkonen, L., Hugenschmidt, T., Berninger, P., Gaidatzis, D., Mohn, F., Artus-Revel, C. G., Zavolan, M., Svoboda, P., and Filipowicz, W. (2008). MicroRNAs control de novo DNA methylation through regulation of transcriptional repressors in mouse embryonic stem cells. *Nat Struct Mol Biol* 15, 259–267.
- Stark, G. R., Kerr, I. M., Williams, B. R., Silverman, R. H., and Schreiber, R. D. (1998). How cells respond to interferons. *Annu Rev Biochem* 67, 227–264.
- Su, H., Trombly, M. I., Chen, J., and Wang, X. (2009). Essential and overlapping functions for mammalian Argonautes in microRNA silencing. *Genes Dev* 23, 304–317.
- Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., and Yamanaka, S. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131, 861–872.
- Takahashi, K., and Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126, 663–676.
- Tam, O. H., Aravin, A. A., Stein, P., Girard, A., Murchison, E. P., Cheloufi, S., Hodges, E., Anger, M., Sachidanandam, R., Schultz, R. M., and Hannon, G. J. (2008). Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 453, 534–538.
- Tay, Y., Zhang, J., Thomson, A. M., Lim, B., and Rigoutsos, I. (2008). MicroRNAs to Nanog, Oct4 and Sox2 coding regions modulate embryonic stem cell differentiation. *Nature* 455, 1124–1128.
- Thomson, J. M., Newman, M., Parker, J. S., Morin-Kensicki, E. M., Wright, T., and Hammond, S. M. (2006). Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes Dev* 20, 2202–2207.
- Ule, J., Jensen, K., Mele, A., and Darnell, R. B. (2005). CLIP: a method for identifying protein-RNA interaction sites in living cells. *Methods* 37, 376–386.
- Ule, J., Jensen, K. B., Ruggiu, M., Mele, A., Ule, A., and Darnell, R. B. (2003). CLIP identifies Nova-regulated RNA networks in the brain. *Science* 302, 1212–1215.

- Vasudevan, S., Tong, Y., and Steitz, J. A. (2007). Switching from repression to activation: microRNAs can up-regulate translation. *Science* 318, 1931–1934.
- Verdel, A., and Moazed, D. (2005). RNAi-directed assembly of heterochromatin in fission yeast. *FEBS Lett* 579, 5872–5878.
- Viswanathan, S. R., Daley, G. Q., and Gregory, R. I. (2008). Selective blockade of microRNA processing by Lin28. *Science* 320, 97–100.
- Wang, Y., Baskerville, S., Shenoy, A., Babiarz, J. E., Baehner, L., and Blelloch, R. (2008). Embryonic stem cell-specific microRNAs regulate the G1-S transition and promote rapid proliferation. *Nat Genet* 40, 1478–1483.
- Wang, Y., Melton, C., Medvid, R., Jaenisch, R., and Blelloch, R. (2007). DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nat Genet* 39, 380–385.
- Warner, D. R., Bhattacharjee, V., Yin, X., Singh, S., Mukhopadhyay, P., Pisano, M. M., and Greene, R. M. (2004). Functional interaction between Smad, CREB binding protein, and p68 RNA helicase. *Biochem Biophys Res Commun* 324, 70–76.
- Watanabe, T., Takeda, A., Tsukiyama, T., Mise, K., Okuno, T., Sasaki, H., Minami, N., and Imai, H. (2006). Identification and characterization of two novel classes of small RNAs in the mouse germline: retrotransposon-derived siRNAs in oocytes and germline small RNAs in testes. *Genes Dev* 20, 1732–1743.
- Watanabe, T., Totoki, Y., Toyoda, A., Kaneda, M., Kuramochi-Miyagawa, S., Obata, Y., Chiba, H., Kohara, Y., Kono, T., and Nakano, T., et al. (2008). Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* 453, 539–543.
- Wianny, F., and Zernicka-Goetz, M. (2000). Specific interference with gene function by double-stranded RNA in early mouse development. *Nat Cell Biol* 2, 70–75.
- Wu, L., and Belasco, J. G. (2005). Micro-RNA regulation of the mammalian lin-28 gene during neuronal differentiation of embryonal carcinoma cells. *Mol Cell Biol* 25, 9198–9208.
- Wulczyn, F. G., Smirnova, L., Rybak, A., Brandt, C., Kwidzinski, E., Ninnemann, O., Strehle, M., Seiler, A., Schumacher, S., and Nitsch, R. (2007). Post-transcriptional regulation of the let-7 microRNA during neural cell specification. *FASEB J* 21, 415–426.
- Yang, N., and Kazazian, H. H. (2006). L1 retrotransposition is suppressed by endogenously encoded small interfering RNAs in human cultured cells. *Nat Struct Mol Biol* 13, 763–771.
- Yang, S., Tutton, S., Pierce, E., and Yoon, K. (2001). Specific double-stranded RNA interference in undifferentiated mouse embryonic stem cells. *Mol Cell Biol* 21, 7807–7816.
- Yi, R., Doehle, B. P., Qin, Y., Macara, I. G., and Cullen, B. R. (2005). Overexpression of exportin 5 enhances RNA interference mediated by short hairpin RNAs and microRNAs. *RNA* 11, 220–226.
- Yi, R., Pasolli, H. A., Landthaler, M., Hafner, M., Ojo, T., Sheridan, R., Sander, C., O’Carroll, D., Stoffel, M., Tuschl, T., and Fuchs, E. (2009). DGCR8-dependent microRNA biogenesis is essential for skin development. *Proc Natl Acad Sci U S A* 106, 498–502.
- Yi, R., Qin, Y., Macara, I. G., and Cullen, B. R. (2003). Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes Dev* 17, 3011–3016.
- Yu, J., Vodyanik, M. A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J. L., Tian, S., Nie, J., Jonsdottir, G. A., Ruotti, V., and Stewart, R., et al. (2007). Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318, 1917–1920.